



Product Datasheet

Product Name	Recombinant Canine Clusterin
Cata No	CB501283
Source	<i>Escherichia Coli.</i>
Synonyms	CLI, AAG4, KUB1, SGP2, SGP-2, SP-40, TRPM2, MGC24903, Glycoprotein 80, Gp80, CLU, Clusterin, Apolipoprotein J, Apo-J.

Description

Clusterin also named Apolipoprotein J (APO-J) is a 75-80 kD disulfide-linked heterodimeric protein containing about 30% of N-linked carbohydrate rich in sialic acid but truncated forms targeted to the nucleus have also been identified.

The precursor polypeptide chain is cleaved proteolytically to remove the 22-mer secretory signal peptide and subsequently between residues 227/228 to generate the a and b chains. These are assembled in anti-parallel to give a heterodimeric molecule in which the cysteine-rich centers are linked by five disulfide bridges and are flanked by two predicted coiled-coil α -helices and three predicted amphipathic α -helices.

Across a broad range of species clusterin shows a high degree of sequence homology ranging from 70% to 80%. It is nearly ubiquitously expressed in most mammalian tissues and can be found in plasma, milk, urine, cerebrospinal fluid and semen. It is able to bind and form complexes with numerous partners such as immunoglobulins, lipids, heparin, bacteria, complement components, paraoxonase, beta amyloid, leptin and others. Clusterin has been ascribed a plethora of functions such as phagocyte recruitment, aggregation induction, complement attack prevention, apoptosis inhibition, membrane remodeling, lipid transport, hormone transport and/or scavenging, matrix metalloproteinase inhibition.

A genuine function of clusterin has not been defined.

One tempting hypothesis says that clusterin is an extracellular chaperone protecting cells from stress induced insults caused by degraded and misfolded protein precipitates.

Clusterin is up- or down regulated on the mRNA or protein level in many pathological and clinically relevant situations including cancer, organ regeneration, infection, Alzheimer disease, retinitis pigmentosa, myocardial infarction, renal tubular damage, autoimmunity and others.

Apolipoprotein-J canine Recombinant produced in E.Coli is a single, non-glycosylated, Polypeptide chain containing 433 amino acids and having a molecular mass of 50.6 kDa.

The protein is fused to His tag at N-Terminus.

The Apolipoprotein-J canine is purified by proprietary chromatographic techniques. The amino acids sequence is identical to UniProtKB/Swiss-Prot entry P25473 amino acids 23–445.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Purity

Greater than 95% as determined by SDS PAGE.

Formulation

Canine Clusterin was lyophilized from PBS, pH 7.5.

*** For Non-Clinical Research Use Only ***



Reconstitution

Add deionized H₂O to a working volume of 0.5mg/ml and let the lyophilized pellet dissolve completely.

Stability

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to **avoid repeated freezing/thawing cycles**. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C. The lyophilized protein remains stable until the expiry date when stored at -20°C.

Sequence

Product Datasheet

MKHHHHHHAS DQAVSDELQ EMSTEGSKYI
NKEIKNALKG VKQIKTLIEQ TNEERKSLLS
NLEEAKKKKE DALNDTKDSE TKLKASQGVG
NDTMMALWEE CKPCLKQTCM KFYARVCRSG
SGLVGHQLEE FLNQSSPFYF WMNGDRIDSL
LENDRQQTHA LDVMQDSFNR ASSIMDELQ
DRFFTREPDQ TYHYSFSLF QRRPFFNPKF
RIARNIIPFP RFQPLNFHDM FQPFFDMIHQ
AQQAMDVNLH RPYHFPIEF PEEDNRTVCK
EIRHNSTGCL KMKDQCEKCQ EILSVDCSSN
NPAQVQLRQE LSNSLQIAEK FTKLYDELLQ
SYQEKMFNTS SLLKQLNEQF SWVSQLANLT
QSEDPFYLQV TTVGSQTSDS NVPVGFTKVV
VKLFSDPIT VMPEAVSRN NPKFMETVAE
KALQEYRQKHREE